

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## MINOR NOTICES

The fresh-water flora of Germany, Austria, and Switzerland.—Paschers has begun the publication of a series of brochures dealing with the fresh-water flora of Germany, Austria, and Switzerland, assisted by numerous specialists. The plan includes 16 parts, 4 of which have just appeared: no. 2 (pp. 192. figs. 398), Flagellatae 2, by A. Pascher (Chrysomonadinae, Crytomonadinae, and Chloromonadineae) and E. Lemmermann (Eugleninae); no. 3 (pp. 66. figs. 69. M 1.80), Dinoflagellatae (Flagellatae 3), by A. J. Schilling; no. 9 (pp. 51. figs. 89. M 1.80), Zygnemales, by O. Borge and A. Pascher; no. 10 (pp. 192. figs. 398. M 4), Bacillariales (Diatomeae), by H. v. Schönfeldt. The numerous illustrations and analytical keys should make the recognition of forms comparatively easy.—J. M. C.

William Russell Dudley.—Leland Stanford Junior University has published a "Dudley Memorial Volume," containing a paper by the late Professor Dudley and appreciations and contributions by friends and colleagues. The list of scientific papers is as follows: "The vitality of Sequoia gigantea," by W. R. Dudley; "The morphology and systematic position of Calycularia radiculosa," by D. H. Campbell; "Studies of irritability in plants. III. The formative influence of light," by G. J. Peirce; "The gymnosperms growing on the grounds of Stanford University," by Leroy Abrams; "The Synchytria in the vicinity of Stanford University," by James McMurphy; "The law of geminate species," by D. S. Jordan; "Some relations between salt plants and salt spots," by W. A. Cannon; "North American species of the genus Amygdalus," by W. F. Wight.—J. M. C.

## NOTES FOR STUDENTS

Cultures of the Uredineae.—The publications of 1912 on cultural work with the plant rusts show an increasing tendency in all parts of the world to clear up by systematic efforts rather than by sporadic cultures the problems of biological relationships in this group of parasitic fungi. In the United States, Arthur, who for many years has been prominently associated with this field of research, reports six species which either have been shown to be autocious or have been connected with their antithetic generation for the first time. These are as follows: Puccinia Lygodesmiae Ellis et Ev. from Lygodesmia juncea (Pursh) D. Don produced teleutospores on the same host without the intercalation of pycnidia or other spore forms. Aecidiospores of Aecidium monoicum Peck from Arabis sp. produced uredinia and telia on

<sup>&</sup>lt;sup>5</sup> PASCHER, A., Die Süsswasser-Flora, Deutschlands, Osterreichs, und der Schweiz. Parts 2, 3, 9, and 10. Jena: Gustav Fischer. 1913.

<sup>6</sup> ARTHUR, J. C., Cultures of Uredineae in 1911. Mycologia 4:49-65. 1912.

<sup>&</sup>lt;sup>7</sup> Unless otherwise stated, teleutosporic material was used in making the infections and the teleutosporic host is given first.